



**University of  
Zurich**<sup>UZH</sup>

**Zurich Open Repository and  
Archive**

University of Zurich  
University Library  
Strickhofstrasse 39  
CH-8057 Zurich  
[www.zora.uzh.ch](http://www.zora.uzh.ch)

---

Year: 2013

---

## **Little-known brachiopods from the Cretaceous of the Helvetic realm of NE Switzerland (Alpstein) and W Austria (Vorarlberg)**

Sulser, Heinz ; Friebe, Georg ; Kürsteiner, Peter

DOI: <https://doi.org/10.1007/s00015-013-0144-6>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-89458>

Journal Article

Published Version

Originally published at:

Sulser, Heinz; Friebe, Georg; Kürsteiner, Peter (2013). Little-known brachiopods from the Cretaceous of the Helvetic realm of NE Switzerland (Alpstein) and W Austria (Vorarlberg). *Swiss Journal of Geosciences*, 106(2):397-408.

DOI: <https://doi.org/10.1007/s00015-013-0144-6>

# Little-known brachiopods from the Cretaceous of the Helvetic realm of NE Switzerland (Alpstein) and W Austria (Vorarlberg)

Heinz Sulser · Georg Friebe · Peter Kürsteiner

Received: 6 September 2012 / Accepted: 10 August 2013 / Published online: 16 November 2013  
© Swiss Geological Society 2013

**Abstract** Three species from different stratigraphical levels of the Cretaceous of the Helvetic Alps are described. (1) Rhynchonellid specimens from the upper Öhrli-Kalk (Öhrli Formation, Late Berriasian) of NE Switzerland (Alpstein) identified as *Lamellaerhynchia heimi* (Sulser 2008) [Rhynchonellida, Hemithiridoidea]. Its range appears to be limited to a small area of the carbonate platform of the northern Alpstein chain. Based on internal and external morphological criteria *L. heimi* differs from other species of *Lamellaerhynchia*, as well as from *Burrirhynchia* cf. *sayni* (Jacob & Fallot 1913), occurring in the younger carbonate platform of the Schrattenkalk Formation (Early Aptian). (2) Recently collected material in various localities of the Altmann Member (Tierwis Formation, Late Hauterivian to Early Barremian) in the Alpstein area identified as *Oblongarcula* cf. *alemannica* Owen 1977 [Terebratulidina, Laqueoidea]. This species is closely related or identical to *O. alemannica* of the North European Boreal province and gives a reference to the occurrence of the genus *Oblongarcula* in the Tethyan domain of the Alps. Partially silicified specimens enable in rare cases a direct access to internal structures after that they were exposed by

an acid treatment. (3) *Tulipina koutaisensis* (Loriol 1896) [Terebratulidina, Kingenoidea], known from Aptian deposits in the central Caucasus of Georgia, has been recorded as a rare species in the Plattenwald-Bed (Selun Member of the Garschella Formation, Albian) of W Austria (Vorarlberg). The localised occurrence and the temporal gap between the Caucasian and the Helvetic *T. koutaisensis* suggest an east–west directed migration along the northern margin of the Tethys Ocean during the Early Cretaceous.

**Keywords** Brachiopoda · Systematics · Early/Middle Cretaceous · Swiss and Austrian Helvetic

## Abbreviations

NMSG Naturmuseum St. Gallen, Switzerland  
MHNG Muséum d'histoire naturelle de la ville de Genève, Switzerland

## 1 Introduction

Recent studies on the Early and Middle Cretaceous brachiopods of the Helvetic Alps are sparse despite the fact that their occurrence has been documented since long by Moesch (1878), Vacek (1879) and by the geological works of Albert Heim (1905), Arnold Heim (1910–1916), Heim and Baumberger (1933) and Heim and Seitz (1934). More recently, two studies about brachiopods of the Cretaceous of Vorarlberg in W Austria were published and included new species (Sulser and Föllmi 1984; Sulser and Friebe 2002), and Sulser (2008) has provided an overview on the present knowledge of the Cretaceous brachiopods in NE Switzerland (Alpstein, Churfirsten) and Vorarlberg (Sulser 2008). This is in contrast to numerous recent publications,

---

Editorial handling: D. A. T. Harper & D. Marty.

---

H. Sulser (✉)  
Paläontologisches Institut und Museum der Universität Zürich,  
Karl Schmid-Strasse 4, 8006 Zurich, Switzerland  
e-mail: heinz.sulser@sunrise.ch

G. Friebe  
Inatura Erlebnis Naturschau, Jahngasse 9,  
6850 Dornbirn, Austria

P. Kürsteiner  
Naturmuseum St. Gallen, Museumstr. 32,  
9000 St. Gallen, Switzerland

which deal with the brachiopods of the same period in non-alpine domains throughout Europe. In the present paper, some little-known species are described which is an important complement to the brachiopod fauna of the alpine Cretaceous.

## 2 Geographical and geological setting

The brachiopods described in this study were collected from six localities in the Alpstein, Switzerland and from one locality in Vorarlberg, Austria (Fig. 1). These are covered by the topographical maps *Landeskarte der Schweiz 1:25,000, Säntis (Blatt 1115)*, and *Diepoldsau (Blatt 1096)*, and by the geological maps *Geologischer Atlas der Schweiz 1:25,000, Säntis (Blatt 1115)*, *Diepoldsau (1096)*, *St. Margrethen (Blatt 1076)*, and *Feldkirch (Blatt 1116)*.

Details regarding the chronostratigraphic stages, formations and other lithostratigraphic units of the above six localities are provided in the descriptions of the individual


brachiopod species. For locality Öhrliattel refer to *Lamellaerhynchia heimi*, for the localities Altmannsattel, Löchlibetter, Tierwis, Chreialp-Litten, Wildhuser Schafboden-Tristen refer to *Oblongarcula cf. alemannica*, and for the locality Plattenwald near Klaus refer to *Tulipina koutaisensis*.

## 3 Materials and methods

The specimens were generally articulated, but frequently crushed or deformed. Even when qualitatively satisfactory, beak, foramen and interarea were often deformed, so that specimens that permitted the examination of both internal and external morphology are rather rare.

The shells are figured in dorsal-, ventral-, lateral- and anterior view. In the anterior view the dorsal valve is at the top.

To study the internal morphology we used methods depending on the nature of preservation. These sections were made each 0.1–0.2 mm, and they were briefly etched



**Fig. 1** Map showing grey-coloured areas of the Säntis nappe of the Alpstein, northeastern Switzerland (**a**) and in Vorarlberg, Western Austria (**b**). Locations where brachiopods were collected are marked by black circles. **a** Locality 1 is Öhrliattel (1.5 km northeast of the Säntis), locality 2 is Altmannsattel (type locality of the Altmann Member, Point 2368, 120 m northwest of the Altmann), locality 3 is Löchlibetter (south of Point 2162, 400 m northeast of the Altmann),

locality 4 is Tierwis (Paratype locality of the Altmann Member, on the path to the Grenzchopf, 1.2 km west of the Säntis), locality 5 is Chreialp-Litten (1.5 km south of Altmann), locality 6 is Wildhuser Schafboden-Tristen (1 km southeast of the Wildhuser Schafberg), and locality. **b** Locality 7 is Plattenwald (near Klaus, 10 km northeast of Feldkirch, Vorarlberg). Black triangles represent mountain peaks

in 5 % hydrochloric acid, rinsed, dried, soaked with acetone and pressed on to acetate-foil. The foils were magnified, and with transmitted light the projections were hand-drawn. The distance from the posterior end of the ventral valve is indicated in millimetres.

For brachiopods in a phosphorite matrix, acetate peels proved mostly unsuitable for the recognition of fine structures. In such cases the sections were photographed. In the case of silicified brachiopods, in addition to the section method, the internal structures could be carefully exposed and directly observed through stepwise application of 2–3 % hydrochloric acid. The mostly incomplete silicification of internal structures was a handicap for achieving satisfactory results, which therefore are rather an exception.

The brachiopods in this study were largely collected by one of us (P. Kürsteiner), mostly together with Karl Tschanz (Zürich). The newly collected material was integrated into the collection of the Naturmuseum St. Gallen (NMSG) and the Muséum d'histoire naturelle de la ville de Genève (MHNG). Additionally, we have also studied specimens in the collections of the Naturmuseum St. Gallen (NMSG), Paläontologisches Institut und Museum der Universität Zürich (PIMUZ), inatura Erlebnis Naturschau Dornbirn (VNS), Muséum d'histoire naturelle de la ville de Genève (MHNG), and the Geological Institute of the Swiss Federal Institute of Technology in Zürich (ETHZH, Churfürsten-collection, Arnold Heim).

#### 4 Systematic palaeontology

Phylum	Brachiopoda Duméril 1806
Subphylum	Rhynchonelliformea Williams et al. 1996
Class	Rhynchonellata Williams et al. 1996
Order	Rhynchonellida Kuhn 1949
Superfamily	Hemithiridoidea Rzhonsnitskaia 1956
Family	Cyclothyrididae Makridin 1955

Subfamily Cyclothyridinae Makridin 1955

Genus *Lamellaerhynchia* Burri 1953

Type species *Terebratula rostriformis* Roemer 1836

*Lamellaerhynchia heimi* (Sulser 2008)

?1916 *Rhynchonella irregularis* (non Pictet); Arn. Heim: p. 459

1999 *Burrirhynchia* sp. Sulser: p. 88 (with Fig.)

2008 *Burrirhynchia heimi* Sulser: p. 103, Fig. 5a–c

*Species name.* In honour of Arnold Heim, who in 1907 defined the Öhrlikopf as the type locality of the Öhrli Formation.

*Holotype.* The specimen described as the holotype (NMSG P. 5189) is 13.8 mm long, 15.3 mm in width and 7.4 mm thick (Fig. 2a) and was recovered from the upper Öhrli-Kalk (Öhrli Formation) of the north face of the Muschelenberg (=type locality of the species). It is depicted in dorsal-, ventral-, lateral-, and frontal view.

*Type locality.* North face of the Muschelenberg, towards Hinter-Öhrlihueb, Kanton Appenzell I.R., Northeastern Switzerland (Fig. 1a).


*Occurrence.* Muschelenberg, Hüenerberg, Hängeten, Altenalpütturm (all from Alpstein).

*Age.* Late Berriasian: Upper Öhrli-Kalk of the Öhrli Formation (Revision by Grasmück-Pflüger 1962).

*Material.* Ten well-preserved specimens (from a total of about 100) from the locality Öhrlisattel (Fig. 1a).

*Diagnosis.* Shell moderately biconvex, shallow-uniplicate, in outline subtrigonal, multicostate, beak straight to sub-erect, pedicle foramen rimmed, septalium and low dorsal septum present, crura raduliform.

*Description.* External morphology (Fig. 2a, b). Shell moderately equibiconvex, on average 14 (8–17) mm long, 16 (8–21) mm wide and 8.5 (4–12) mm thick, when more



**Fig. 2** a–b *Lamellaerhynchia heimi* (Sulser), Upper Öhrli-Kalk (Öhrli Formation), both specimens from Öhrlisattel. **a** Holotype (NMSG P. 5189); **b** paratype (NMSG P. 5190), dorsal, ventral, lateral and anterior view. **c–d** *Burrirhynchia* cf. *sayni* (Jacob & Fallot),

Rawil Member (Schrattenkalk Formation), both specimens from Mattstock (Bärenfall) near Amden (Canton St. Gallen, Switzerland), dorsal, ventral and anterior view (NMSG P. 5191). Scale bar 1 cm


than 10 mm long, width always slightly wider than long. Bluntly trigonal outline, anterior commissure uniplicate, rarely almost rectimarginate. Dorsal valve without fold, ventral valve with weakly developed sulcus anteriorly. About 20 (14–24) radial costae, covering the whole shell, rounded in profile, rarely bifurcating. Beak prominent, straight to suberect, pedicle foramen rimmed, round to oval, hypothyrid. Deltidial plates fused. Beak ridges with blunted edges, apical angle  $90^{\circ}$ – $105^{\circ}$ .

Internal morphology (Fig. 3). Ventral valve proximally with slightly converging, distally with subparallel dental plates, toward the plane of articulation detached of the inner wall of the valve. Dorsal valve without cardinal process, septalium present (section 3.0), but reduced. Hinge plates proximally connected by a thin, roof-shaped lamella (sections 3.2 and 3.6), then separated and aligning horizontally (section 4.3). Dental sockets deeply rounded, semi-circular in cross section, in which hinge teeth equipped with a dentalium are inserted obliquely (section 4.3). Crural bases reinforced, crura raduliform, dorsally slightly concave, with jagged ends (section 6.2). Median septum low, thick (section 3.2), persisting up to one third of the valve length.

**Discussion.** This species, introduced by Sulser (2008) and only briefly commented, can be assigned to the Cyclothyridae on account of the rimmed foramen, the prominent beak, and the shape of the crura. The determining skeletal features for the reassignment from the originally ascribed genus *Burrirhynchia* to *Lamellaerhynchia* are the shell covered with radial costae, the wedge-shaped dorsal


septum, and the reduced septalium. In comparison to *L. heimi*, *L. rostriformis* (Roemer 1836) is larger, the outline is more wide-rounded, the rib-pattern coarser, the individual ribs more angular in profile and the septalium is missing. The areal region of *L. rostriformis* is more distinctive, the interarea flatter, more extensive (cf. Burri 1956, Owen and Thurell 1968). *L. renauxiana* (d'Orbigny 1850) grows into a significantly larger and comparatively thicker form. Additionally it shows a pronounced uniplication, which is partially asymmetrically distorted. Otherwise, *L. heimi* keeps rather constant in its shape. It exhibits discrete characteristics, which distinguish it as a separate species.

The establishment of *L. heimi* as a separate species, however, requires further clarification. The species from the Öhrli-Kalk has been mixed up with the small rhynchonellid *Burrirhynchia gibbsiana* var. *sayni* (Jacob & Fallot 1913) from the Early Aptian Schrattenkalk Formation (Rawil Member, previously lower Orbitolinen-Bed). The confusion of the two species was partially due to the fact that early alpine geologists were not able to strictly distinguish the lithologically similar Öhrli- and Schrattenkalk. *B. gibbsiana* var. *sayni* occurs throughout the Helvetic realm (it has often been referred to *Rhynchonella gibbsiana* or *R. gibbsi*, from which the obsolete name “Gibbsi-beds” was derived). Owen (1956) had shown that the *Burrirhynchia gibbsiana* (J. de C. Sowerby 1826) occurs in the Early Albian of southern England as an endemic species. The form from the Helvetic Alps, provisionally named here *Burrirhynchia* cf. *sayni*, was originally described from the Barremian of the French Alps. Its internal structure is unknown.



**Fig. 3** *Lamellaerhynchia heimi*, Upper Öhrli-Kalk (Öhrli Formation), from Öhrlisattel. Transverse serial sections, specimen 16.4 mm long, 17.4 mm wide and 10.2 mm thick. The numbers of the sections indicate the distance in mm from the posterior end of the shell. Scale bar 1 cm





**Fig. 4** *Burrirhynchia* cf. *sayni*, Rawil Member (Schrattenkalk Formation), from Mattstock (Bärenfall) near Amden (Canton St.Gallen, Switzerland). Transverse serial sections, specimen 13.0 mm long, 15.2 mm wide and 11.3 mm thick. Scale bar is 1 cm

*Lamellaerhynchia heimi* and *Burrirhynchia* cf. *sayni* differ both in external shape and in internal structure. *L. heimi* is slimmer and less globular. *B. cf. sayni* has finer costae, their number is higher (25–35), the anterior commissure wide uniplicate, the shell generally less wide (Fig. 2c, d). In *B. cf. sayni* septalium and dorsal septum are missing, the strong dental plates progress horizontally, the crura are shorter, extending to about one third valve length (Fig. 4).

At the type locality, the occurrence of *L. heimi* seems to be restricted to a discrete horizon in the lower part of the upper Öhrli-Kalk where it is abundant and associated with fragments of the oyster *Alectryonia* (*Lopha*). The horizon in question was referred to by Grasmück-Pflüger (1962) as a gnarly-schistous, oncolithic, soft band of marlstone containing echinoderm remains and sponge spicules. *L. heimi* seems to show a certain degree of frequency variability in its lateral persistence, which could indicate a facies dependency. In shallow-water carbonates, characterised by reef forming corals, *L. heimi* is associated with the brachiopods *Psilothyris tamarindus* (J. de C. Sowerby) and *Loriolithyris valdensis* (Loriol). While the occurrence of *L. heimi* is limited to the type area of the Helvetic region, *L. valdensis* has a wider geographic and stratigraphic distribution.

*Lamellaerhynchia heimi* can readily be distinguished from other cyclothyrid rhynchonellids and can be regarded as the ancestral form of the genus. It is the characteristic fossil of the upper Öhrli-Kalk, and it remains unclear whether it occurs in other, facially and temporally comparable deposits of the Cretaceous Helvetic or elsewhere. With regard to relevant log descriptions, which document


the fossil assemblages this seems unlikely (Burger 1985). The equivalent of the eastern continuation of the Öhrli Formation in Vorarlberg is the shallow-water facies of the Örfli Formation. This contains a different brachiopod assemblage, not examined in detail as yet, but where *L. heimi* seems to be absent. Föllmi et al. (2007) regard these two stratigraphic units as synonymous and the younger name (Örfli) as obsolete.

*Lamellaerhynchia heimi* of Barremian and *Burrirhynchia* cf. *sayni* of Aptian age occur in slightly marly beds of two carbonate platforms with a large temporal gap of some tens of millions of years. The external resemblance might result from an analogous mode of life in similar palaeo-environments. A similar case can be observed in the terebratulids *Loriolithyris valdensis* of the Öhrlikalk and *Selliathyris sella* (J. de C. Sowerby) of the Schrattenkalk. Whether the morphological resemblance is due to adaptations in two coral dominated platforms needs to be substantiated by examining more examples. A point which has to be paid attention to also is the polymorphism and polytypism of both species.

Order	Terebratulida Waagen 1883
Suborder	Terebratellidina Muir-Wood 1955
Superfamily	Laqueoidea Thomson 1927
Family	Terebrataliidae Richardson 1975
Subfamily	Gemmarculinae Elliott 1947
Genus	<i>Oblongarcula</i> Elliott 1959

Type species *Terebratula oblonga* J. de C. Sowerby 1829

*Oblongarcula* cf. *alemannica* Owen 1977



**Fig. 5** a–c *Oblongarcula* cf. *alemannica* Owen, Altmann Member (Tierwis Formation). **a** From Löchlibetter (NMSG coll. Kürsteiner 5A.09.05), dorsal, ventral, lateral and anterior view. **b** From Altmannsattel, size series showing variation of outline and costation (NMSG coll. Kürsteiner 5A.06.18). **c** From Altmannsattel (NMSG coll. Kürsteiner 5A.06.18), anterior view showing zigzag pattern of

costae (= same specimen as the left one of Fig. 5b). **d** *Oblongarcula alemannica* Owen, Hauterivian/Barremian, from Mönchewahlberg near Wolfenbüttel, about 10 km south of Braunschweig, Lower Saxony, Germany, dorsal, ventral, lateral and anterior view (NMSG P. 5192). Scale bar is 1 cm

1836 *Terebratula oblonga* Sow.; Roemer: 46, Pl. 2, Fig. 23  
 1838 *Terebratula oblonga* Sow.; von Buch: 159, Pl. 16, Fig. 2

1839 *Terebratula pectiniformis* Roemer (non Schlotheim): 20, Pl. 18, Fig. 9

1848 *Terebratula oblonga* d'Orbigny (sic!): 113, Pl. 515, Fig. 7–19

1850 *Terebratula oblonga* Sow.; Strombeck: 76, Pl. 4, Fig. 1–19

1864 *Terebratula oblonga* Sow.; Meyer: 254, Pl. 11, Fig. 12–14

1871 *Terebratula puscheana* Roemer; Quenstedt: 275, Pl. 44, Fig. 139

1874 *Terebratula oblonga* Sow.; Davidson: 26, Pl. 2, Fig. 29–31

1878 *Waldheimia oblonga* Sow.; Moesch: 34

1916 *Rhynchonella lata* (non d'Orb.); Heim: 398

1933 *Terebratella oblonga* d'Orb. (!); Heim & Baumberger: 175, 180

? 1933 *Terebratella neocomiensis* (non d'Orb.); Heim & Baumberger: 176

1959 *Oblongarcula oblonga* (J. de C. Sowerby); Elliott: 147

1977 *Oblongarcula alemannica* Owen: 227–230, Textfig. 14, Pl. 3, Fig. 5

1999 *Burrirhynchia?* sp. Sulser: p. 88–89 (with Fig.)

2008 *Oblongarcula* cf. *alemannica* Owen; Sulser: 106, Abb. 5, Fig. Q–U

**Occurrence.** Northwestern Germany; northeastern France; northeastern Switzerland: Alpstein (Altmannsattel, Chreialp, Löchlibetter, Tierwis, Wildhuser Schafboden), Churfirsten (Glatthalde [doubtful locality description by

Arnold Heim]); Austria: Vorarlberg (Hohenems, Unterklien, Breiterberg).


**Age.** Late Hauterivian/Early Barremian; Altmann-Member of the Tierwis Formation.

**Material.** 25 well preserved specimens from different localities of the Altmann region (Fig. 1a).

**Description.** External morphology (Fig. 5a–c). Shell acutely biconvex, rectimarginate, subpentagonal to elongate-oval or bluntly trigonal in outline (Fig. 5b), on average 17.5 (6.5–26) mm long, 12 (5.5–20) mm wide and 9.5 (3.0–20) mm thick. 10–25 strong, deeply incised, occasionally bifurcating and intercalating costae, which draw a zigzag-line at the anterior commissure (Fig. 5c). Beak straight to suberect, beak-ridges sharp. Interarea flat, hinge line curved. Pedicle foramen large, round, mesothyrid, deltidial plates fused. Apical angle about 60°.


**Internal morphology** (Figs. 6, 7). Ventral valve: dental lamellae lightly concave, hinge teeth blunt (Fig. 7b, section 1.4; Fig. 7c, section 1.6). Dorsal valve: cardinal process poorly developed if at all (see black point in Fig. 7b, section 2.4 and Fig. 7c, section 1.6). Dental sockets with well-marked inner and outer ridges. Septalium present (Fig. 7a, sections 2.8 and 3.0), hinge plates fused, becoming weakly inclined and triangular in connection with median septum (Fig. 7c, sections 2.2–3.0). Septum persists through mid-valve. Crural bases in juvenile specimen with pointed ends of descending branches (Fig. 6b). Loop not observed.

**Discussion.** The shell is distinguished by marked ribs and might be misidentified as a rhynchonellid brachiopod when the calcitic shell structure is silicified and cannot be



**Fig. 6 a, b** *Oblongarca* cf. *alemannica* Owen, Altmann Member (Tierwis Formation). Silicified internal structures prepared by exposure to acid. **a** Dorsal valve interior from Tierwis near Grenzchopf, showing with compact hinge plate (NMSG P. 5193). **b** Juvenile specimen from Löchlibetter, showing a pair of dental lamellae with


lateral umbonal cavities, in between initial stages of articulated hinge plates (*horizontal arrow*), crural bases with pointed ends of descending branches (*vertical arrow*), in the center a knob, probably the initial stage of median septum (*oblique arrow*) (NMSG P. 5194). Scale bars 5 mm



**Fig. 7 a–c** *Oblongarca* cf. *alemannica* Owen, Altmann Member (Tierwis Formation), Transverse serial sections. **a** Specimen 17.6 mm long, 12.4 mm wide and 9.0 mm thick, from Löchlibetter. **b** Specimen 18.1 mm long, 13.7 mm wide and 11.1 mm thick, from Löchlibetter;

**c** Specimen 12.7 mm long, 10.2 mm wide and 8.3 mm thick, from Altmannsattel. The numbers of the sections indicate the distance in mm from the posterior end of the shell. Scale bars are 1 cm





**Fig. 8** *Oblongarcula alemannica* Owen, Hauterivian/Barremian, from Mönchewahlberg near Wolfenbüttel, about 10 km south of Braunschweig, Lower Saxony, Germany. Transverse serial sections,

specimen 18.7 mm long, 12.5 mm wide and 12.8 mm thick. The numbers of the sections indicate the distance in mm from the posterior end of the shell. Scale bar is 1 cm

assessed. Kempf (1966, p. 12) who logged the section at the Altmannsattel referred to these brachiopods in question as “Rhynchonellen”. Sulser (1999) also referred to them as a questionable *Burrirhynchia?* sp., before more favourably preserved specimens were found.

The serial sections show these brachiopods in an entirely different light. An important diagnostic feature of the brachidium is a plane-undivided, slightly trough-shaped valve plate, which is supported by a high, drawn-out dorsal septum. Partially silicified specimens, after acid preparation permitted direct observation of this characteristic element of the interior organisation (Fig. 6a). This strongly supports their assignment to the terebratulid genus *Oblongarcula*. The genera *Psilothyris* Cooper and *Belothyris* Smirnova show a similar internal organisation, but their smooth shells exclude a possible confusion. The affirmation for the assignment finally resulted from the comparison with the serial sections of *Oblongarcula alemannica* published by Owen (1977).

*Oblongarcula* has long been known from the Hilskonglomerat (Hauterivian) in northwestern Germany (Schöppenstedt and Berklingen near Braunschweig) and northeastern France. Roemer (1836) referred to it as *Terebratula oblonga*, which J. de C. Sowerby (1829) had established for correspondent forms in South England and which has become the type species of *Oblongarcula* (Elliott 1959). Owen (1977) compared the German and English specimens and established the new species *O. alemannica* for the former. We figure a German specimen from Mönchewahlberg near Braunschweig (Fig. 5d) and have sectioned another one (Fig. 8). The separation into two species is hardly visible in the exterior morphology and relies essentially on a minimal divergence of the internal structure. This is a blunt, short protuberance located in the centre of the proximal hinge plate (Owen 1977, text-fig. 14, sections 6–9), which cannot be recognized in *O. oblonga* (Owen 1977, text-fig. 16). Owen himself qualified his approach by regarding *O. alemannica*

as a possible regional variation of *O. oblonga*. The short temporal interval between *O. alemannica* (Hauterivian) and *O. oblonga* (Aptian) can also not justify a separation of both taxa. Therefore, the status of two independent species is still questionable. Nevertheless, we adhere to *O. alemannica* and take it as reference species by naming the relevant specimens from the Helvetic Altmann Member *Oblongarcula* cf. *alemannica*.

Our collected material occurs in a glauconitic, schistous limestone, near the base of the Altmann Member, partially enriched in a shellbed and associated with echinoid spines. Besides elongate shells, especially among the smaller examples, wide and flat forms can be distinguished, which possess length/width-ratios near 1:1. In contrast, the variability of the rib density- and refinement seems to be independent of individual size (Fig. 5b, showing individuals in the size series).

Superfamily Kingenoidea Elliott 1948  
Family Kingenidae Elliott 1948  
Subfamily Kingeninae Elliott 1948  
Genus *Tulipina* Smirnova 1962

Type species *Zeilleria koutaisensis* de Loriol 1896

*Tulipina koutaisensis* (de Loriol 1896)


1896 *Zeilleria Koutaisensis* Loriol: 145–147; Pl. 5, Fig. 19–24  
1962 *Tulipina koutaisensis* (Loriol); Smirnova: 103–105; Textfig. 5, 6

1972 *Tulipina koutaisensis* (Loriol); Smirnova: 104–105; Pl. 9, Fig. 7

1990 *Tulipina koutaisensis* (Loriol); Smirnova: 136–137, Textfig. 85, Pl. 35, Fig. 2, 3

2002 *Kingena*-like, unidentified brachiopods: Sulser & Friebe: 423, Fig. 13A, B

Type locality. Kutaissi (Georgia) at the south foot of the Central Caucasus Mountains, about 60 km east of the Black Sea.



**Fig. 9 a, b** *Tulipina koutaisensis* (Loriol). **a** Plattenwald Member (Garschella Formation, Albian), from Plattenwald, Vorarlberg, Austria (NMSG P. 5195), dorsal, lateral and ventral view. **b** Late

Hauterivian to Early Barremian, from Koutaisi, Central Caucasus, Georgia (MHNG GEPI 81032), dorsal, ventral, lateral and anterior view. Scale bars 1 cm

**Occurrence.** Koutaisi (Georgia), Plattenwald near Klaus (Vorarlberg), Austria (Fig. 1b).

**Age.** Late Barremian to Early Aptian of Georgia; Albian of Vorarlberg (Garschella Formation: Plattenwald-Bed of the Selun Member).

**Material.** 2 specimens from the locality Plattenwald in Vorarlberg (Fig. 1g)

**Description.** External morphology (Fig. 9a). Shell small (11 mm long, 10 mm wide and 9 mm thick, only 2 specimens inspected), smooth, strongly biconvex, globose with subcircular outline, anterior commissure flat to slightly sulcate, lateral commissure straight. Beak thick, much incurved, pressed towards the posterior part of dorsal valve. Pedicle foramen small, apical region rounded.

Internal morphology (Fig. 10). Ventral valve: dental lamellae short, not persistent (sections 1.7–2.8). Dorsal valve: dental sockets and articulation of hinge teeth diffuse (sections 4.0 and 4.3). Cardinal process not observed. Septalium deep, cup-like (sections 3.6–4.0). Crural bases reinforced (section 4.5). Median septum proximately high, distally becoming low and truncated (sections 5.3 and 7.6), persisting through at least mid-valve. Long V-shaped, reflected loop with a pair of broad, obliquely connected ascending bands, descending bands united with median septum (section 5.7).


**Discussion.** The brachiopod fauna from the Plattenwald-Bed of Vorarlberg is dominated by the terebratulid *Moutonithyris dutempleana* (d'Orbigny), whereas the rhynchonellid *Burrirhynchia tripartita* (Pictet) and *Orbirhynchia parkinsoni* (Owen) are not frequent. *Tulipina koutaisensis* is so far only represented by a few specimens. Because of its hood-shaped shell it was provisionally referred to as “kingenoid” in Sulser and Friebe (2002, p. 425, Fig. 13). More extensive studies have now permitted its determination and assignment to *Zeilleria koutaisensis* Loriol, on which Smirnova (1962) based her genus *Tulipina*. The exterior appearance already suggests

that it is identical with the form of the Plattenwald locality (Fig. 9a, b), and the comparison of serial sections with those on a specimen from Favre's (1875) Caucasian collection confirms this (Figs. 10, 11). The latter sections are practically identical with the ones in Smirnova (1962, text-fig. 6). In the reconstruction of the internal structures by Smirnova (1962, text-fig. 5) *T. koutaisensis* shows a *Kingenia*-like, long, vertically lying loop, in which the median septum is connected with the descending lamellae.

It is interesting to note that Favre (1875) conducted in 1871 a scientific expedition to the Central Caucasus. The brachiopod-population of more than 20 specimens, collected in Koutaisi—in what today is Georgia—was named *Zeilleria koutaisensis* by de Loriol (1896), and is stored in the Muséum d'histoire naturelle de Genève.


An unusual structural element at the base of the dorsal septum in the sections of the Caucasian specimens (Smirnova 1962, text-fig. 6; Fig. 11, marked by arrow in section 5.2) is missing in the specimens from Plattenwald. These are preserved as moulds and slightly weathered along the septum, on account of which the mentioned structure was probably destroyed. Its function is not known. Maybe it served as an expanded attachment place for the adductor muscle on the dorsal side. With regard to the rather weak seeming anchoring of the valves (Fig. 10, sections 3.6–4.3 and Fig. 11, sections 2.2–2.7), a strengthened adductor muscle could have supported the articulation of these small shells.

Previously *T. koutaisensis* had only been recognised in the type area of the central Caucasus, for which Smirnova (1962) specified a Late Barremian to Early Aptian age. Favre (1875) had already suggested a pre-Albian age based on the co-incidence of ammonites from the group “*Crioceras*” (= *Crioceratites*). Additional verification of the stratigraphical situation is complicated because recent publications from Georgia are unavailable for political reasons. However for the sediments in question, an Albian age can be excluded on the basis of co-occurring ammonites (D. Ruban, Rostow, Ukraine, personal communication).



**Fig. 10** *Tulipina koutaisensis* (Loriol). Plattenwald Member (Garschella Formation, Albian), from Plattenwald near Klaus, Vorarlberg, Austria. Transverse serial sections, specimen 11 mm long, 10.0 mm

wide and 9.1 mm thick. The numbers of the sections indicate the distance in mm from the posterior end of the shell. Scale bar is 1 cm



**Fig. 11** *Tulipina koutaisensis* (Loriol). Late Hauterivian to Early Barremian, from Kuttaisi, Central Caucasus, Georgia. Transverse serial sections, specimen 10.0 mm long 9.3 mm wide and 7.6 mm thick. Scale bar is 1 cm

## 5 Conclusions and prospects

### 5.1 *Lamellaerhynchia heimi*

Within the Berriasian brachiopod assemblage of Europe, the rhynchonellids represent a relatively small group with only a few genera. In her review, Gaspard (1999) mentioned *Lacunosella*, *Monticlairella* und *Peregrinella* as the main genera. At present, the Berriasian *Lamellaerhynchia heimi* is the oldest alpine representative of the genus. From the Valanginian onwards it is widespread among Early Cretaceous neritic brachiopods of Europe, northern Africa, Mexico and western Russia with about 20 described species. *Lamellaerhynchia* is well represented by distinct

species in the Jura Mountains of northwestern Switzerland and probably occurs with further species in the Cretaceous of the Helvetic realm.

### 5.2 *Oblongarcula*

Its occurrence in the alpine Cretaceous confirms the occasional occurrence of *Terebratella oblonga* in fossil lists of geological works (Moesch 1878; Heim and Baumberger 1933). This genus has so far been described from the Boreal province of northern Europe and southern England. Some questionable species, probably belonging to *Oblongarcula* have been described, such as *Terebratella neocomiensis* d'Orbigny and *T. exquisita* Loriol, see Pictet

(1872). It seems that *Oblongarcula* advanced as far as the Helvetic realm in the border zone to the Tethys. Assuming that a transgression occurred in the latest Hauterivian (Haq et al. 1987), faunal exchanges between the individual zones have become possible.

### 5.3 *Oblongarcula* cf. *alemannica*

Silicified specimens can be prepared by treatment with acid and directly reveal details of the internal structure, but complete silicification of all calcitic parts is rare. However, as shells of small and medium-sized specimens are common in the collected material, there is a potential to document early stages (see Fig. 6b and its comment) and reconstruct ontogenetical development, usually only known from recent brachiopods. This is a difficult work, but should be carried out in the future, because it offers the chance to elucidate internal structures being built in the course of individual growth in extinct genera such as *Oblongarcula* (see e.g. Gaspard 2003).

### 5.4 *Tulipina koutaisensis*

In a comparative study of Early Cretaceous brachiopods, Lobacheva (1986) stated that many “European” species are known from an earlier age in western and central Asia. This also holds true for *Tulipina koutaisensis*. This species had previously been recorded from the Late Barremian to Early Aptian of the central Caucasus in Georgia. The time gap until it appeared in the strongly condensed Early to Late Albian (?Early Cenomanian) Plattenwald-Bed would comprise about 10 million years. Covering—what is today—an about 2,500 km long distance corresponds to a migration rate of 25 m per 100 years. Sessile brachiopods can only actively move during a short phase of the larval stage. The rate of dispersal is influenced by several factors, which makes it difficult to estimate. One of the factors is the prevailing westerly ocean-current along the northern margin of the Tethys, which transported *T. koutaisensis* from Georgia to Vorarlberg, the only two places from where the species is presently known. Föllmi (1986, 1989) reported the unusual occurrence of a dropstone in the Albian of Vorarlberg, which was probably derived from gneiss in the Bohemian Massif, and he ascribed its eastern provenance to the prevailing ocean current at that time. *T. koutaisensis* in the Helvetic of Vorarlberg would at least support this assumption.

### 5.5 General conclusions

The Early Cretaceous brachiopods of Europe to a certain extent show both widely-distributed and strictly endemic species. Palaeogeographical and palaeoecological factors

influence this pattern, but also the autecology of the brachiopods and, therefore, their resistance or adaptability towards such factors. Taxonomical uncertainties are still a handicap in answering the question to what extent an “alpine group” of various stratigraphical levels can be distinguished from groups of other provenances. In the light of these questions an improved taxonomical record of the involved brachiopod species in realms such as the Helvetic Alps remains an actual task.

**Acknowledgments** We wish to express our gratitude for the opportunity to study the collections of the Geological Institute of the Swiss Federal Institute of Technology Zürich (Milena Pika-Biolzi) and inatura Erlebnis Naturschau Dornbirn, Vorarlberg. We would especially like to acknowledge Lionel Cavin (Muséum d’histoire naturelle Genève), who kindly supplied us with brachiopods from the Cretaceous of the Caucasus Mountains (collection of E. Favre) both for comparison and preparation. We wish to thank Dmitri Ruban (Rostov, Ukraine) for information on the geological and stratigraphical context of the relevant brachiopods from the Caucasus, and Karl Föllmi (Université Lausanne) for clarification of palaeogeographical questions regarding the Cretaceous of the Helvetic realm. We also thank Richard Waite (Ithaca NY, USA) for the English translation of the original German manuscript and Joëlle Soom (Heimiswil) for the map with the relevant settings of sampled brachiopods. We greatly appreciate the reviews of Danièle Gaspard (Université Paris-Sud, France) and Michael Sandy (University of Dayton OH, USA) whose proposals and critical comments helped to improve the manuscript.

## References

- Buch, L. von (1838). Essai d’une classification et d’une description des Térébratules. Mémoire de la Société géologique de France 3, 105–238, Pl. 13–20.
- Burger, H. (1985). Palfries-Formation, Öhrli-Formation und Vitznau-Mergel (Basale Kreide des Helvetikums) zwischen Reuss und Rhein. Stratigraphische, fazielle, mineralogische und paläogeographische Untersuchungen. Mitteilungen aus dem Geologischen Institut der Eidg. Technischen Hochschule und der Universität Zürich, Neue Folge Nr. 254.
- Burri, F. (1956). Die Rhynchonelliden der Unteren Kreide (Valanginien-Barrémien) im westschweizerischen Juragebirge. *Eclogae Geologicae Helveticae*, 49, 599–701, 15 Pl.
- Davidson, T. (1874). A monograph of the fossil Brachiopoda. 4/1. Supplement to the recent, tertiary and cretaceous species (pp. 1–72). London: Palaeontographical Society (Monogr.), 8 Pl.
- Elliott, G. (1959). Six new genera of Mesozoic Brachiopoda. *Geological Magazine*, 96, 146–148.
- Favre, E. (1875). Recherches géologiques dans la partie centrale de la Chaîne du Caucase. *Denkschriften der schweizerischen naturforschenden Gesellschaft*, 27, 1–91.
- Föllmi, K.B. (1986). Die Garschella- und Seewer Kalk-Formation (Aptian-Santonian) im Vorarlberger Helvetikum und Ultrahelvetikum. Mitteilungen des Geologischen Instituts der ETH und der Universität Zürich, Neue Folge Nr. 262.
- Föllmi, K.B. (1989). Evolution of the Mid-Cretaceous Triad. Lecture Notes in Earth Sciences (Vol. 23). Berlin: Springer.
- Föllmi, K. B., Bodin, S., Godet, A., Linder, P., & van de Schootbrugge, B. (2007). Unlocking paleo-environmental information from Early Cretaceous shelf sediments in the Helvetic



- Alps: Stratigraphy is the key! *Swiss Journal of Geosciences*, 100, 349–369.
- Gaspard, D. (1999). Distribution of Lower Cretaceous brachiopods in Europe (Berriasian to Aptian). *Bulletin de la Société géologique de la France*, 170, 311–326.
- Gaspard, D. (2003). Some Cretaceous long-looped terebratulide brachiopods analysed in the light of the diversity observed in the ontogeny of recent representatives. *Bulletin de la Société géologique de la France*, 174, 261–269.
- Grasmück-Pflüger, M. (1962). Mikrofazielle Beobachtungen an den Öhrschichten (Berriasian) der Typuslokalität. *Eclogae Geologicae Helveticae*, 55, 417–442.
- Haq, B. U., Hardenbol, J., & Vail, P. R. (1987). Chronology of fluctuating sea levels since the Triassic. *Science*, 235, 1156–1167.
- Heim, Arn. (1905). II. Teil: Westlicher Teil des Säntisgebirges. In A. Heim (Ed.), Das Säntisgebirge. Textband. *Beiträge zur geologischen Karte der Schweiz* [N.F.], 16.
- Heim, Arn. (1907). Gliederung und Fazies der Berrias-Valanginian-Sedimente in den helvetischen Alpen. *Vierteljahrsschrift der Naturforschenden Gesellschaft Zürich*, 52, 484–499.
- Heim, A. (1910/1913/1916). Monographie der Churfürsten-Mattstock-Gruppe. I–III. Teil. Beiträge zur geologischen Karte der Schweiz [N.F.] 20, 1–272, 16 Pl.(1910); 20, 273–368, 8 Pl.(1913); 20, 369–574, 8 Pl. (1916).
- Heim, Am., & Baumberger, E. (1933). Jura und Unterkreide in den helvetischen Alpen beiderseits des Rheins (Vorarlberg und Ostschweiz). *Denkschriften der Schweizerischen naturforschenden Gesellschaft*, 68, 155–220, 2 Pl.
- Heim, A., & Seitz, O. (1934). Die mittlere Kreide in den helvetischen Alpen von Rheintal und Vorarlberg und das Problem der Kondensation. *Denkschriften der Schweizerischen naturforschenden Gesellschaft* 69, 185–310, 3 Pl.
- Jacob, Ch., & Fallot, P. (1913). Etude sur les Rhynchonelles portlandiennes, néocomiennes et mésocrétacées du Sud-Est de la France. *Abhandlungen der Schweizerischen Paläontologischen Gesellschaft* 39, 11 Pl.
- Kempf, T.A. (1966). Geologie des westlichen Säntisgebirges. *Beiträge zur geologischen Karte der Schweiz* [N.F.], 128, 2 Pl.
- Lobacheva, S.V. (1986). Early Cretaceous brachiopods of the South USSR and some new data on Capillithyridinae. In P.R. Racheboeuf & C.C. Emig (Eds.), *Les Brachiopodes fossils et actuels. Biostratigraphie du Paléozoïque* (Vol. 4, pp. 129–134).
- Loriol, P. de. (1896). Note sur quelques brachiopodes crétacés recueillis par M.E. Favre dans la chaîne centrale du Caucase et dans le Néocomien de la Crimée. *Revue Suisse de la Zoologie et Annales du Musée d'Histoire naturelle de Genève* 4, pp. 135–163, 4 Pl.
- Meyer, C. J. A. (1864). Notes on the Brachiopoda from the Pebblebed of the Lower Greensand of Surry; with descriptions of the new species, and remarks on the correlation of the Greensand Beds of Kent, Surrey and Berks, and the Farrington Sponge-gravel, and the Tourtia of Belgium. *Geological Magazine*, 1, 249–257, Pl. 11–12.
- Moesch, C. (1878). Zur Paläontologie des Säntisgebirges. Über neue und weniger bekannte Petrefacten aus der Kreide des Sentisgebirges. In A. Escher v.d. Linth (Ed.), *Geologische Beschreibung der Sentis-Gruppe. Beiträge zur geologischen Karte der Schweiz* (Vol. 13), 3 Pl.
- Orbigny, A. d' (1847/1851). Paléontologie française. Description des mollusques et rayonnés fossiles. Terrains crétacés. 4: Brachiopodes (1847), Pl. 490–599 (1851).
- Owen, E.F. (1956). The Lower Cretaceous brachiopods '*Rhynchonella*' gibbsiana (J. de C. Sowerby) and *Sulcirhynchia hythensis*, sp. nov. *Annual Magazine Natural History* [12], 9, 164–172, 1 Pl.
- Owen, E.F. (1977). Evolutionary trends in some Mesozoic Terebratellacea. *Bulletin of the British Museum (Natural History), Geology*, 28, 205–253, 3 Pl.
- Owen, E.F., & Thurell, R.G. (1968). British Neocomian rhynchonellid brachiopods. *Bulletin of the British Museum (Natural History)*, 16, 99–123, 4 Taf.
- Pictet, F.J. (1872). In F.J. Pictet & G. Campiche (Eds.), Description des fossiles du terrain crétacé des environs de Sainte-Croix. 5. partie: Brachiopodes (terminé par P. de Loriol). Matériaux de la Paléontologie Suisse, 14 Pl.
- Quenstedt, F.A. (1871). Petrefactenkunde Deutschlands. Tübingen und Leipzig. 1. Abt. 2. Band. Die Brachiopoden: 1–160 (1868); 161–464 (1869); 465–748 (1870); 25 Taf. (1871).
- Roemer, F.A. (1836). Die Versteinerungen des norddeutschen Hannover: Oolithen-Gebirges, 16 Pl.
- Roemer, F.A. (1839). Die Versteinerungen des norddeutschen Oolithen-Gebirges. Hannover: Ein Nachtrag, 5 Pl.
- Smirnova, T. N. (1962). New data on Lower Cretaceous Dallinids (Brachiopods). *Paleontological Zhurnal*, 1962, 97–105. (in russian).
- Smirnova, T.N. (1972). Early Cretaceous brachiopods of the Crimea and the northern Caucasus. *Akademia Nauk SSSR (Moskwa)*, 13 Pl. (in Russian).
- Smirnova, T. N. (1990). Systematik der Kreide-Brachiopoden. *Akademia Nauk SSSR (Moskwa)*, 40 Pl. (in Russian).
- Sowerby, J. de C. (1826–1829). The Mineral Conchology of Great Britan 6, Pl. 504–609.
- Strombeck, E. (1850). *Terebratula oblonga*. Briefliche Mitteilung an Herrn L. von Buch. *Zeitschrift der deutschen geologischen Gesellschaft*, 2, 76–82, 1 Pl.
- Sulser, H. (1999). *Die fossilen Brachiopoden der Schweiz und der angrenzenden Gebiete*. Juragebirge und Alpen: Paläontologisches Institut und Museum der Universität Zürich.
- Sulser, H. (2008). Die Brachiopoden aus der alpinen Kreide der Nordostschweiz (Alpstein, Churfürsten, Mattstock) und von Vorarlberg—ein Überblick. *Berichte der St.Gallischen Naturwissenschaftlichen Gesellschaft* 91, pp. 97–122, 4 Pl.
- Sulser, H., & Föllmi, K. B. (1984). Eine neue Brachiopodenart (*Lacunosella acutifrons* n.sp., Rhynchonellida) aus dem helvetischen « Gault » Vorarlbergs (Österreich). *Eclogae Geologicae Helveticae*, 77, 619–629.
- Sulser, H., & Friebe, J. G. (2002). Brachiopods from the Plattenwald Bed (Albian, Cretaceous) of the Helvetic Alps of Vorarlberg (Austria). *Eclogae Geologicae Helveticae*, 95, 415–427.
- Vacek, M. (1879). Über Vorarlberger Kreide. Eine Localstudie. *Jahrbuch der kaiserlich-königlichen geologischen Reichsanstalt* 29, 659–756, 3 Pl.